



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

PROVE that, if $0 < \alpha < \beta$,

$$\int_{\alpha}^{\beta} \log \frac{\beta - x}{x - \alpha} \frac{dx}{x} = \frac{1}{2} \left[\log \frac{\beta}{\alpha} \right]^{2*}.$$

[Frank Morley.]

SOLUTION.

The integration is best effected by use of the following theorem, or lemma .

$$\int_{\alpha}^{\beta} \varphi \left[\frac{x}{\beta} \right] \frac{dx}{x} = \int_{\alpha}^{\beta} \varphi \left[\frac{a}{x} \right] \frac{dx}{x},$$

where $\varphi(z)$ is any finite and continuous function of z ; and β and α have any fixed values, provided when one is zero the other must be infinite.

To prove this, let $\frac{x}{\beta} = \frac{a}{y}$; whence

$$\frac{dx}{x} = - \frac{dy}{y}.$$

$$\therefore \int_{\alpha}^{\beta} \varphi \left[\frac{x}{\beta} \right] \frac{dx}{x} = - \int_{\beta}^{\alpha} \varphi \left[\frac{a}{y} \right] \frac{dy}{y} = \int_{\alpha}^{\beta} \varphi \left[\frac{a}{x} \right] \frac{dx}{x}.$$

Now

$$I \equiv \int_{\alpha}^{\beta} \log \frac{\beta - x}{x - \alpha} \frac{dx}{x} = \int_{\alpha}^{\beta} \log \left\{ \frac{\beta \left[1 - \frac{x}{\beta} \right]}{x \left[1 - \frac{\alpha}{x} \right]} \right\} \frac{dx}{x};$$

whence

$$I = \int_{\alpha}^{\beta} \log \frac{\beta}{x} \frac{dx}{x} + \int_{\alpha}^{\beta} \log \left[1 - \frac{x}{\beta} \right] \frac{dx}{x} - \int_{\alpha}^{\beta} \log \left[1 - \frac{\alpha}{x} \right] \frac{dx}{x}.$$

By the above lemma, the last two integrals are equal. We have, also,

$$\frac{dx}{x} = - d \log \frac{\beta}{x};$$

whence

$$I = - \int_{\alpha}^{\beta} \log \frac{\beta}{x} d \left[\log \frac{\beta}{x} \right] = \frac{1}{2} \left[\log^2 \frac{\beta}{x} \right]_{\beta}^{\alpha},$$

* The index, 2, is omitted in the text; it is undoubtedly an error on the part of the printer.

and

$$\int_a^{\beta} \log \frac{\beta - x}{x - a} \frac{dx}{x} = \frac{1}{2} \left[\log \frac{\beta}{a} \right]^2. \quad [H. L. Rice.]$$

380

THE sides of a variable rectangle pass through four fixed points. Find the position of the rectangle and its dimensions when its area is a maximum.

[Geo. R. Dean.]

SOLUTION.

Let A, B, C, D be the four points in order; a, b the diagonals AC, BD ; and α the angle between them. Then if φ be inclination of a side p to AC , the two sides are

$$p = a \cos \varphi, \quad q = b \sin (\varphi + \alpha);$$

and the area

$$pq = ab \sin (\varphi + \alpha) \cos \varphi = \frac{1}{2} ab [\sin (2\varphi + \alpha) + \sin \alpha]$$

is a maximum when $\varphi = 45^\circ - \frac{1}{2} \alpha$. The sides are

$$\frac{1}{2} a \sqrt{2} (\sin \frac{1}{2} \alpha + \cos \frac{1}{2} \alpha), \quad \frac{1}{2} b \sqrt{2} (\sin \frac{1}{2} \alpha + \cos \frac{1}{2} \alpha),$$

and are equally inclined to the bisectrices of the angles between AC and BD .

[Wm. M. Thornton.]

Solved also by H. L. Rice and G. B. M. Zerr.

381

FROM a point in the circumference of a circle of radius R as centre is described the external arc of a circle of radius r . Determine r so that the area of the lune shall equal that of the original circle. [W. M. Thornton.]

SOLUTION.

Let φ be the angle at the center of the R -circle which subtends the chord

$$r = 2R \sin \frac{1}{2} \varphi.$$

Equating the area of the lune to that of the R -circle, we have

$$\frac{1}{2} r^2 (\pi + \varphi) - R^2 \varphi + rR \cos \frac{1}{2} \varphi = \pi R^2,$$

whence

$$\tan \varphi = \pi + \varphi.$$

From which the value of φ , determined by trial, is

$$\varphi = 77^\circ 27' 12''.08$$

Whence

$$r = 1.25122 R. \quad [W. H. Echols.]$$

Solved also by G. B. M. Zerr and H. L. Rice.

382

FOUR equal circles tangent to each other cut off equal areas from a given circle. Required the radii of the cutting circles when the aggregate area cut off from the given circle is the greatest possible. [*Artemas Martin.*]

SOLUTION.

The distance between the centres of the fixed circle and cutting circle $= r\sqrt{2}$, where r is the radius of a cutting circle. Let R = radius of given circle. Then

$$4r^2 \cos^{-1} \left[\frac{3r^2 - R^2}{2\sqrt{2}r^2} \right] + 4R^2 \cos^{-1} \left[\frac{R^2 + r^2}{2\sqrt{2}Rr} \right] - 2\sqrt{6R^2r^2 - R^4 - r^4}$$

is the area cut off by the four circles. Differentiating and reducing, we get

$$r^2 \cos^{-1} \left[\frac{3r^2 - R^2}{2\sqrt{2}r^2} \right] = \frac{1}{2} \sqrt{6R^2r^2 - R^4 - r^4}. \quad (1)$$

Let

$$\cos \theta = \frac{3r^2 - R^2}{2\sqrt{2}r^2}. \quad (2)$$

Whence (1) becomes $\theta = \sqrt{2} \sin \theta$, the solution of which gives $\theta = 79^\circ 43' 46''$. From (2),

$$r = .633 R.$$

(1) is also satisfied by $r = (\sqrt{2} + 1) R$, and $r = (\sqrt{2} - 1) R$. The first of these values gives the four tangent circles circumscribed to the given circle, the second, the four tangent circles inscribed in the given circle.

[*G. B. M. Zerr.*]

383

If c' , c'' , c''' be the sides of any triangle inscribed in an ellipse, and b' , b'' , b''' the semi-diameters parallel to the sides, show that the area is

$$A = abc'c''c''' / (4b'b''b'''). \quad [W. O. Whitescarver.]$$

SOLUTION I.

It is shown by Salmon (Conic Sections, p. 220) that if the triangle be given by the eccentric angles α , β , γ , its area is

$$= 2ab \sin \frac{1}{2}(\alpha - \beta) \sin \frac{1}{2}(\beta - \gamma) \sin \frac{1}{2}(\gamma - \alpha).$$

He also shows on page 219 that

$$c' = 2b' \sin \frac{1}{2}(\alpha - \beta), \quad c'' = 2b'' \sin \frac{1}{2}(\beta - \gamma), \quad c''' = 2b''' \sin \frac{1}{2}(\gamma - \alpha);$$

whence by simple substitution we have the result above.

SOLUTION II.

The radius of the circumscribing circle is $\frac{b'b''b'''}{ab}$ (Salmon, p. 220); whence, at once,

$$A = \frac{abc'c''c'''}{4b'b''b'''} . \quad [W. O. Whitescarver.]$$

Solved also by G. B. M. Zerr, H. L. Rice, and F. G. Radelfinger.

384

If c be a chord of an ellipse through the points whose eccentric angles are α and β , and b' the semi-diameter parallel to the chord, show that the area of the triangle formed by the chord and the tangents at its extremities is

$$S = \frac{abc^2}{4b'^2} \tan \frac{1}{2} (\alpha - \beta) . \quad [W. O. Whitescarver.]$$

SOLUTION.

The equations of the two tangents are

$$\frac{x \cos \alpha}{a} + \frac{y \sin \alpha}{b} = 1 ,$$

$$\frac{x \cos \beta}{a} + \frac{y \sin \beta}{b} = 1 .$$

Their intersection has for its coordinates

$$x = \frac{a \cos \frac{1}{2} (\alpha + \beta)}{\cos \frac{1}{2} (\alpha - \beta)} , \quad y = \frac{b \sin \frac{1}{2} (\alpha + \beta)}{\cos \frac{1}{2} (\alpha - \beta)} .$$

$$\therefore \frac{2S}{ab} = \begin{vmatrix} \cos \alpha & \sin \alpha & 1 \\ \cos \beta & \sin \beta & 1 \\ \frac{\cos \frac{1}{2} (\alpha + \beta)}{\cos \frac{1}{2} (\alpha - \beta)} & \frac{\sin \frac{1}{2} (\alpha + \beta)}{\cos \frac{1}{2} (\alpha - \beta)} & 1 \end{vmatrix} ,$$

or

$$\begin{aligned} \frac{S}{ab} &= -\sin \frac{1}{2} (\alpha - \beta) \cos \frac{1}{2} (\alpha - \beta) + \cos^2 \frac{1}{2} (\alpha + \beta) \tan \frac{1}{2} (\alpha - \beta) \\ &\quad + \sin^2 \frac{1}{2} (\alpha + \beta) \tan \frac{1}{2} (\alpha - \beta) . \end{aligned}$$

$$\therefore S = ab \tan \frac{1}{2} (\alpha - \beta) \sin^2 \frac{1}{2} (\alpha - \beta) ,$$

or, by Exercise 383,

$$S = \frac{abc^2}{4b'^2} \tan \frac{1}{2} (\alpha - \beta) . \quad [H. L. Rice.]$$

Solved also by W. O. Whitescarver, F. G. Radelfinger, and G. B. M. Zerr.

385

If in exercise 384 a tangent be drawn parallel to the chord, show that the base of the triangle formed will be $c \sec \frac{1}{2} (a - \beta)$, and its area $ab \tan^3 \frac{1}{2} (a - \beta)$. [W. O. Whitescarver.]

SOLUTION.

If the base be t , we see that it touches the ellipse at the point $\gamma = \frac{1}{2} (a + \beta)$ (see Salmon, Conic Sections, p. 219). Therefore Δ the area of triangle is, from Exercise 373, by substituting for γ ,

$$\Delta = ab \tan \frac{1}{2} (a - \beta) \tan^2 \frac{1}{4} (a - \beta) \quad (1)^*$$

or

$$= 2ab \frac{\tan^3 \frac{1}{4} (a - \beta)}{1 - \tan^2 \frac{1}{4} (a - \beta)}.$$

Since this triangle and that of Ex. 384 are similar we have $\frac{\Delta}{S} = \frac{t^2}{c^2}$, which gives

$$t = 2b' \tan \frac{1}{4} (a - \beta). \quad (2)^*$$

[W. O. Whitescarver.]

Solved also by H. L. Rice and G. B. M. Zerr.

392

A SYSTEM of great circles intersect upon the equator of a sphere; a curve is drawn connecting points on the spherical surface where the circles of this system make a constant angle α with the meridians. Show that the stereographic projection of any such curve is a circular cubic whose equation may be written

$$\tan \alpha \tan \theta = \frac{c^2 + r^2}{c^2 - r^2},$$

c being the radius of the sphere, and one of the poles being the centre of the projection. [R. A. Harris.]

SOLUTION.

In stereographic projection the distance of any point from the origin is equal to the radius of the sphere multiplied by the tangent of one-half the polar distance, or $r = c \tan \frac{1}{2} \varphi$; whence

$$\sec \varphi = \frac{r^2 + c^2}{r^2 - c^2}.$$

If α is the constant angle, and θ the complement of the angle between the given meridian and the meridian of the moving point, we have

$$\sec \varphi = \tan \alpha \tan \theta. \quad [\text{Geo. R. Dean.}]$$

* The values given in the problem are wrong. The error came by mistaking $\frac{1}{4}$ for $\frac{1}{2}$ in equation (1). This of course changed (2).